


We are unable to supply this entire article because the publisher requires payment of a copyright fee. You may be able to obtain a copy from your local library, or from various commercial document delivery services.

From Forest Nursery Notes, Winter 2012

**184. © Survival and early growth of mixed forest stands installed in a Mediterranean region: effects of site preparation intensity.** Fonseca, F., de Figueiredo, T., and Martins, A. *Forest Ecology and Management* 262:1905-1912. 2011.



## Survival and early growth of mixed forest stands installed in a Mediterranean Region: Effects of site preparation intensity

Felícia Fonseca<sup>a,\*</sup>, Tomás de Figueiredo<sup>a,1</sup>, Afonso Martins<sup>b,2</sup>

<sup>a</sup> Centro de Investigação de Montanha (CIMO), Escola Superior Agrária de Bragança, Campus de Sta Apolónia, Apartado 1172, 5301-855 Bragança, Portugal

<sup>b</sup> Universidade de Trás-os-Montes e Alto Douro, Apartado 1013, 5001-911 Vila Real, Portugal

### ARTICLE INFO

#### Article history:

Available online 20 February 2011

#### Keywords:

Mediterranean Region  
*Pseudotsuga menziesii*  
*Castanea sativa*  
 Plantation  
 Mortality  
 Growth

### ABSTRACT

In Mediterranean environments, availability of water and nutrients are the main factors limiting the success of afforestation. As part of a wider project, an experiment was established in Northeast Portugal, aiming at testing the effect of several site preparation techniques on plant survival and growth (height and diameter) in a newly installed mixed forest stand. Results presented regard plant response during 42 months after plantation. The experimental protocol consisted in seven treatments described by mechanical operations that rank soil disturbance intensity from none to high, set in plots of 375 m<sup>2</sup>, randomly distributed in three blocks, in different topographic positions (gentle slope plateau, moderate slope shoulder, and steep mid-slope). *Pseudotsuga menziesii* (PM) and *Castanea sativa* (CS) forest species were planted in a 4 m × 2 m scheme and in alternate rows with 12 plants on each row per plot, summing up 72 plant per specie and treatment at start of the experiment. The results show that: (i) the highest mortality was observed immediately after the plantation and before the dry season, on the lowest intensity treatments; (ii) after the dry season, the highest mortality was also observed in treatments with the lowest intensity of soil disturbance, while the lowest values were found on the intermediate intensity treatments; (iii) during the experimental period, the effect of treatments on plant growth (height and diameter) was statistically significant; however, experimental results do not lead yet to a clear quantitative relationship between soil disturbance intensity due to site preparation and plant response under the conditions tested.

© 2011 Elsevier B.V. All rights reserved.

### 1. Introduction

In Mediterranean environments, availability of water and nutrients are the main factors limiting the success of afforestation (Daget, 1977; Rey, 1998; Ojasvi et al., 1999; Bocio et al., 2004). Afforestation programmes in Portugal have to take into account these soil and climatic constraints, as Mediterranean climate prevails in most of the territory (Ribeiro, 1986; Costa et al., 1998). In Portugal the success and productivity of most forest plantations is limited by poor soil conditions, namely a low root support capacity, which has negative effects on the amount of available water and nutrients. Soil preparation operations are therefore required before planting, using more or less intense tillage in order to increase soil depth, as well as water and nutrient availability, and, so, improving soil conditions for plant growth (Worrell and Hampson, 1997; Fisher and Binkley, 2000; Querejeta et al., 2001).

Several studies have been made on the effect of surface soil tillage using scarification and herbicide application for weeds and shrub control and improving root depth (McLaughlin et al., 2000; Archibold et al., 2000; Burgess and Wetzel, 2000; Wetzel and Burgess, 2001). However, there are only few references on the effect of deep tillage on soil properties and plant response (Fernandes and Fernandes, 1998; Fisher and Binkley, 2000; Querejeta et al., 2001; Martins and Pinto, 2004; Carlson et al., 2006). New studies are therefore necessary to improve our knowledge and to support decisions on best operation selection according to site conditions. The wide diversity of mechanical site preparation techniques that may be applied emphasizes the need for studies on newly installed forest stands, especially in areas where information is still limited, as it is particularly the case of the Mediterranean Region (Varelides and Kritikos, 1995). Furthermore, most studies are performed in adult stands, and so the installation phase is less understood, often lacking important components of the initial dynamics of these systems (Canham, 1989; Lieberman et al., 1989).

This paper aims at presenting and discussing data collected 42 months after plantation in an experiment carried out to study mortality and growth (height and diameter) of a mixed stand of

\* Corresponding author. Tel.: +351 273303253/273303200; fax: +351 273325405.

E-mail addresses: [ffonseca@ipb.pt](mailto:ffonseca@ipb.pt) (F. Fonseca), [amartins@utad.pt](mailto:amartins@utad.pt) (A. Martins).

<sup>1</sup> Tel.: +351 273 303 200; fax: +351 273 325 405.

<sup>2</sup> Tel.: +351 259 35020; fax: +351 259 350480.