

In: Murphy, P. M. (compiler). The challenge of producing
 native plants for the Intermountain area; Proceedings: Intermountain
 Nurseryman's Association 1983 conference; 1983 August 8-11;
 Las Vegas, NV. Gov. Tech. Rept. INT-168. USDA For. Ser.
 Intermountain For. & Range Exp. Sta. 1984. pp 39-43.

BIOLOGY AND MANAGEMENT OF BOTRYTIS BLIGHT

Robert L. James

ABSTRACT

Botrytis cinerea is an important pathogen of conifer seedlings in western North America, especially within greenhouses. Environmental conditions in greenhouses, such as high humidity and cool temperatures, are conducive to infection by and spread of this fungus. To reduce losses from Botrytis blight, cultural practices aimed at reducing inoculum and altering environmental conditions necessary for infection should be combined with rotated use of different fungicides. Several fungicides used to control Botrytis in the past are no longer effective because the fungus has developed tolerance to them. Fungicides commonly used to control this disease are discussed.

INTRODUCTION

Grey mold caused by *Botrytis cinerea* (Fr.) Pers. is one of the most damaging diseases of seedlings in forest tree nurseries. The disease is especially severe on containerized conifers in greenhouses where conditions are ideal for infection by and buildup of the fungus (James, Woo and Myers 1982; McCain 1978). However, Botrytis blight may also occur in seedbeds where it causes damage during cool, wet portions of the year (James 1980; James and others 1983). The fungus is also responsible for losses to seedlings in storage (Smith and others 1973).

Although many conifer species are susceptible to Botrytis, greatest damage has been reported on Douglas-fir, western hemlock, lodgepole pine, and spruce in British Columbia (Sutherland and Van Eerden 1980), western larch, lodgepole pine, and Engelmann spruce in northern Idaho and northwestern Montana (James and Genz 1983; James and Gilligan 1983; James and others 1982), lodgepole pine, Scots pine, Engelmann spruce and blue spruce in Colorado (Gillman and James 1980), and giant sequoia and Douglas-fir in California (McCain and Smith 1978).

BIOLOGY

Of the 22 species of Botrytis, *B. cinerea*, the one that affects conifer seedlings, is the most common and has the widest host range (over 200 plant species) (Jarvis 1980b; Sutherland and Van Eerden 1980). Other Botrytis species are more pathogenically specialized and thus have narrower host ranges.

Robert L. James, Plant Pathologist, Cooperative Forestry and Pest Management, Northern Region, USDA Forest Service, Missoula, Montana.

A typical disease cycle for *B. cinerea* is shown in figure 1. Initial infection in nurseries results from spores produced on nearby infected plants or crop debris and from fungal resting structures (sclerotia) (Coley-Smith 1980; McCain 1978). Sclerotia often form after the growth phase of the fungus or following seedling mortality (Coley-Smith 1980). These sclerotia persist in soil, plant debris or within greenhouses and can produce both sexual (ascospores) and asexual (conidia) spores.

The sexual stage of the fungus is *Botryotinia fuckeliana* (DeBary) Whetzel, which has been found frequently in nature (Jarvis 1980b). Apothecia produced from overwintering sclerotia give rise to ascospores which may initiate infection (Jarvis 1980a). However, asexual conidia are responsible for most infection, spread, and buildup of the disease in nurseries.




Figure 1.--Disease cycle of *Botrytis cinerea* (after Jarvis 1980a).

Conidia are dry and usually dispersed by air currents and less frequently carried by water droplets (Jarvis 1980a). Conidial dispersal occurs primarily when the relative humidity is rising or falling rapidly (Jarvis 1980a). Presence of free moisture on foliage for several hours and prolonged cool temperatures of about 13-14° C are necessary for infection (Blakeman 1980). Germinating conidia form appressoria on the surface of leaves and germ tubes penetrate directly through the cuticle (Blakeman 1980). Wounded or necrotic

host tissues are quickly infected and colonized (Sutherland and Van Eerden 1980).

Within the disease cycle, the fungus may become inactive (latent) following conidial dispersal or infection (figure 1). However, when inoculum is abundant and environmental and host susceptibility conditions are conducive, "aggressive pathogenicity" occurs (Jarvis 1980a). Conducive environmental conditions include high relative humidity, cool temperatures, and free surface moisture on foliage. Host susceptibility factors include nutrient imbalances causing seedling stress and presence of senescent tissues for saprophytic buildup of inoculum (Sutherland and Van Eerden 1980). When conditions for infection are ideal and inoculum abundant, latent periods are short and epidemics can occur quickly (Jarvis 1980a).

Symptoms of *Botrytis* infection usually become apparent when crowns of conifer seedlings begin to close and affected seedlings usually occur in isolated pockets (Gillman and James 1980; James and others 1982). The fungus usually first attacks senescent tissues at the base of seedlings and then spreads to surrounding live host material (Smith and others 1973; Sutherland and Van Eerden 1980). Symptoms on infected seedlings include needle necrosis, twig and stem lesions, and mortality.

MANAGEMENT

Controlling *Botrytis* blight is difficult because the pathogen is capable of attacking all plant parts at almost any stage of their growth and in storage (Maude 1980). The best approach to control is to avoid conditions that are suited for disease buildup. This includes controlling stocking by reducing density to improve air circulation among seedlings (Cooley 1981), which means producing fewer trees per unit area. However, this is compensated by higher quality, disease-free seedlings. If possible, irrigation during periods of host susceptibility should also be limited (Cooley 1981). Adding drying agents to irrigation water to expedite drying of foliage may also help reduce infection. Fertilization should also be done properly. For example, too much fertilizer may cause seedlings to burn, providing ideal infection courts for *Botrytis* (Sutherland and Van Eerden 1980), and too little fertilizer may stress seedlings making them more susceptible to infection (Cooley 1981). Another important cultural practice to reduce loss from *Botrytis* blight is sanitation, aimed primarily at reducing inoculum. Sanitation practices include periodic removal of infected plants and plant debris, and cleaning greenhouse benches and floors with a surface sterilant between crops (Cooley 1981). Potential inoculum sources outside greenhouses, especially those upwind, should be eliminated when possible.

As containerized production of conifers has increased, *Botrytis* blight has become more important. As a result, many growers have had to rely on fungicides to keep losses at acceptable levels. Several fungicides either used operationally or showing promise for future use are listed in table 1. Some

of the more important of these are discussed below.

Benomyl is a systemic fungicide that has been used operationally since the early 1970's. When it was first introduced, benomyl provided excellent control of many diseases over a wide range of crop plants. As a result, many growers began to use it exclusively to control *Botrytis* blight, especially in greenhouses (McCain 1978; Miller and Fletcher 1974). However, as early as 1971 tolerance to benomyl by *Botrytis* was evident (Bollen and Scholten 1971). Since then, there have been many reports of tolerance to this fungicide by different pathogens on a variety of crops including ornamental flowers, vegetables, fruit crops, and conifer seedlings (Cooley 1981; Gillman and James 1980; James and Gilligan 1983; Jarvis and Hargreaves 1973; Miller and Fletcher 1974). Simple tests have been developed to quickly assay presence of tolerant fungal strains. These involve growing the test organisms on nutrient media amended with the fungicide. Such tests have been used to evaluate tolerance of *Botrytis* strains to benomyl and other fungicides throughout the West. Results indicate that tolerance of *Botrytis* to benomyl is so widespread that this chemical is usually ineffective and no longer recommended for use in most nurseries (Cooley 1981; Gillman and James 1980; James and Gilligan 1983).

Chlorothalonil is another fungicide that has been commonly used to control *Botrytis* in greenhouses. However, its ability to adequately control the disease has often been reduced, especially after continued use (James and Gilligan 1983). Recent tests indicate that some *Botrytis* populations in Oregon, Montana, and Colorado are tolerant to chlorothalonil (Cooley 1981; Gillman and James 1980; James and Gilligan 1983). Although tolerance to chlorothalonil is not as widespread as with benomyl, it is fairly common and has been shown to develop quickly in greenhouses (James and Gilligan 1983).

Captan is a general protective fungicide that is fairly effective against *Botrytis* (James and others 1982). However, tolerant strains to this fungicide have also been shown to exist (Cooley 1981; Gillman and James 1980; James and Gilligan 1983; Parry and Wood 1959).

Dicloran is an effective fungicide against *Botrytis* diseases (James and others 1982), even though tolerance of natural *Botrytis* strains has been found (Cooley 1981; Gillman and James 1980; James and Gilligan 1983; Webster and others 1970). Tolerant strains of the fungus can also easily develop in the laboratory (James, unpublished). Therefore, dicloran should not be used repeatedly unless rotated with other fungicides.

Two relatively new fungicides should also be mentioned. Iprodione was originally developed for turf diseases (Danneberger and Vargas 1982; Sanders and others 1978) and shows strong toxicity towards *Botrytis* (Pappas and Fisher 1979; Powell 1982). Vinclozolin is a chemical with specific action against *Botrytis* and related fungi (Pappas and Fisher 1979; Ritchie 1982). Iprodione has

been tested against Botrytis blight of conifers and shows excellent promise (James and others 1982). Vinclozolin was also tested, but showed extensive phytotoxicity to western larch seedlings at label rates (James and Genz 1983). Both fungicides require more field tests and need to be registered for use on conifers. Previous tests (Cooley 1981; James and Gilligan 1983; Leroux and others 1977; Pappas and others 1979) indicate that strains of Botrytis tolerant to iprodione and vinclozolin exists, although not in large numbers. Tolerant strains can also develop rapidly to these fungicides in the laboratory (James, unpublished).

Apparently none of the fungicides currently available can be considered completely effective against all Botrytis strains likely to be encountered. As a result, fungicide usage should be

limited to the minimum amounts necessary for effective disease control. Also, different fungicides should be used in rotation so as not to exert selective pressure on Botrytis populations to develop tolerance. Rotated fungicides should have different modes of action, i.e. systemic chemicals alternated with broad spectrum protectants (Cooley 1981; James and Gilligan 1983).

For effective control of Botrytis blight, cultural practices, such as better sanitation, providing adequate air circulation, and reducing irrigation, should be combined with rotated use of different fungicides. Cultural practices can reduce fungal inoculum and alter environmental conditions necessary for infection, whereas fungicides can protect susceptible plant tissues from infection. The combination of both procedures is necessary for an effective control strategy.

Table 1.--Fungicides used to control Botrytis blight in containerized conifer nurseries.

Fungicide	Trade names	Manufacturers	Chemical name
benomyl	Benlate [®] Tersan 1991 [®] Benomyl	Dupont Lilly Miller	Methyl-1-(butylcarbamoyl)-2 benzimidazole carbamate
captan	Captan Orthocide [®]	Stautier Chevron	N-(Trichloromethyl)thio]-4-cyclohexene-1, 2-dicarboximide
chlorothalonil	Bravo 500 [®] Daconil 2787 [®]	Diamond Shamrock	Tetrachloroisophthalonitrile
copper	Tri-Basic [®]	GP Chemical Phelps-Dodge Cities Service	Basic copper sulfate
dicloran	Botran [®]	Tuco	2,6-Dichloro-4-nitroaniline
ferbam	Carbamate	Dupont	ferric dimethyldithiocarbamate
iprodione	Chipco 26019 [®] Rovral [®]	Rhone-Poulenc	3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboximide
mancozeb	Fore [®]	Dupont	Contains 16% maganese, 2% zinc and 62% ethylenebisdithiocarbamate ion/maganese ethylenebisdithiocarbamate plus zinc ion.
maneb	Dithane M-45 [®]	Rhom & Haas	maganese ethylene bisdithiocarbamate
thiophanate-methyl	Zyban [®]	Mallinckrodt	dimethyl[(1,2-phenylene)bis(iminocarbonothioyl)]bis(carbamate)
thiram	Thylate [®]	Dupont	Tetramethylthiuram disulfide
vinclozolin	Ronilan [®] Ornalin [®]	BASF Mallinckrodt	3-(3,5-dichlorophenyl)-5-ethenyl-5-methyl-2,4-oxazolidinedione
zineb	Zineb Dithane 278 [®]	Rhom & Haas	zinc ethylenebisdithio-carbamate

PUBLICATIONS CITED

- Blakeman, J. P. Behaviour of conidia on aerial plant surface. In: Coley-Smith, J. R.; Verhoeff, K.; Jarvis, W. R., eds. The biology of Botrytis. London: Academic Press; 1980: 115-151.
- Bollen, G. J.; Scholten, G. Acquired resistance to benomyl and some other systemic fungicides in a strain of Botrytis cinerea in cyclamen. Neth. J. Pl. Path. 77: 80-90. 1971.
- Coley-Smith, J.R. Sclerotia and other structures of survival. In: Coley-Smith, J. R.; Verhoeff, K.; Jarvis, W. R., eds. The biology of Botrytis. London: Academic Press; 1980: 85-114.
- Cooley, S. J. Fungicide tolerance of Botrytis cinerea isolates from conifer seedlings. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; 1981. 13 pp.
- Danneberger, T. K.; Vargas, J. M., Jr. Systemic activity of iprodione in Poa annua and post-infection activity for Drechslera sorokiniana leaf spot management. Plant Disease 66(10): 914-915. 1982.
- Gillman, L. S.; James, R. L. Fungicidal tolerance of Botrytis within Colorado greenhouses. Tree Planters' Notes 31(1): 25-28. 1980.
- James, R. L. Engelmann spruce needle blight at the Coeur d'Alene nursery, Idaho. Rept. 80-21. Missoula, MT: U. S. Department of Agriculture, Forest Service, Northern Region; 1980. 5 pp.
- James, R. L.; Genz, D. Fungicide tests to control Botrytis blight of containerized western larch at the Champion Timberlands Nursery, Plains, Montana. Rept. 83-12. Missoula, MT: U. S. Department of Agriculture, Forest Service, Northern Region; 1983. 7 pp.
- James, R. L.; Gilligan, C. J. Fungicidal tolerance of Botrytis cinerea from the Flathead Indian Reservation greenhouse, Ronan, Montana. Rept. 83-5. Missoula, MT: U. S. Department of Agriculture, Forest Service, Northern Region; 1983. 15 pp.
- James, R. L.; Woo, J. Y.; Malone, P. Evaluation of fungicides to control Botrytis blight in western larch seedbeds at the Coeur d'Alene nursery, Idaho. Rept. 83-6. Missoula, MT: U. S. Department of Agriculture, Forest Service, Northern Region; 1983. 8 pp.
- James, R. L.; Woo, J. Y.; Myers, J. F. Evaluation of fungicides to control Botrytis blight of containerized western larch and lodgepole pine at the Coeur d'Alene nursery, Idaho. Rept. 82-17. Missoula, MT: U. S. Department of Agriculture, Forest Service, Northern Region; 1982. 13 pp.
- Jarvis, W. R. Epidemiology. In: Coley-Smith, J. R.; Verhoeff, K.; Jarvis, W. R., eds. The biology of Botrytis. London: Academic Press; 1980a: 219-250.
- Jarvis, W. R. Taxonomy. In: Coley-Smith, J. R.; Verhoeff, K.; Jarvis, W. R., eds. The biology of Botrytis. London: Academic Press; 1980b: 1-18.
- Jarvis, W. R.; Hargreaves, A. J. Tolerance to benomyl in Botrytis cinerea and Penicillium corymbiferum. Plant Pathology 22: 139-141. 1973.
- Leroux, P.; Fritz, R.; Gredt, M. Etudes en laboratoire de souches de Botrytis cinerea Pers., résistantes á la dichlozoline, au dichloran, au quinitozene, á la vinchlozoline et au 26019 RP (ou Glycophene). Phytopathol. Z. 89: 347-348. 1977.
- Maude, R. B. Disease control. In: Coley-Smith, J. R.; Verhoeff, K.; Jarvis, W. R., eds. The biology of Botrytis. London: Academic Press; 1980: 275-308.
- McCain, A. H. Nursery disease problems - containerized nurseries. In: Gustafson, R. W., ed. Western forest nursery council and Inter-mountain Nurseryman's Association: conference and workshop proceedings; Eureka, CA; 1978: B139-142.
- McCain, A. H.; Smith, P. C. Evaluation of fungicides for control of Botrytis blight of container-grown redwood seedlings. Tree Planters' Notes 29(4): 12-13. 1978.
- Miller, M. W.; Fletcher, J. T. Benomyl tolerance in Botrytis cinerea isolates from glasshouse crops. Trans. Br. Mycol. Soc. 62(1): 99-103. 1974.
- Pappas, A. C.; Cooke, B. K.; Jordan, V. W. L. Insensitivity of Botrytis cinerea to iprodione, procymidone and vinclozolin and their uptake by the fungus. Plant Pathology 28(1): 71-76. 1979.
- Pappas, A. C.; Fisher, D. J. A comparison of the mechanisms of actions of vinclozolin, procymidone, iprodione and prochloraz against Botrytis cinerea. Pesticide Science 10: 239-246. 1979.
- Parry, K. E.; Wood, R. K. S. The adaptation of fungi to fungicides: adaptation to captan. Ann. Appl. Biology 47(1): 1-9. 1959.
- Powell, C. C. New chemicals for managing disease on glasshouse ornamentals. Plant Disease 66(2): 171. 1982.
- Ritchie, D. F. Effect of dichloran, iprodione, procymidone, and vinclozolin on the mycelia growth, sporulation, and isolation of resistant strains of Monilinia fructicola. Plant Disease 66(6): 484-486. 1982.

- Sanders, P. L.; Burpee, L. L.; Cole, H., Jr.; Duich, J. M. Control of fungal pathogens of turf grass with the experimental iprodione fungicide, R. P. 26019. Plant Dis. Repr. 62(6): 549-553. 1978.
- Smith, R. S., Jr.; McCain, A. H.; Srago, M. D. Control of Botrytis storage rot of giant sequoia seedlings. Plant Dis. Repr. 57(1): 67-69. 1973.
- Sutherland, J. R.; E. Van Eerden. Diseases and insect pests in British Columbia forest nurseries. Joint Rept. No. 12. British Columbia Ministry of Forests, Canadian Forestry Service; 1980. 55 pp.
- Webster, R. K.; Ogawa, J. M.; Bose, E. Tolerance of *Botrytis cinerea* to 2,6-Dichloro-4-nitro-aniline. Phytopathology 60(10): 1489-1492. 1970.

In: Murphy, Patrick M., compiler. The challenge of producing native plants for the Intermountain area: proceedings: Intermountain Nurseryman's Association 1983 conference; 1983 August 8-11; Las Vegas, NV. General Technical Report INT-168. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station; 1984. 96 p.